



Pergamon

Tetrahedron Letters 40 (1999) 667-670

TETRAHEDRON
LETTERS

Synthesis of an Inositol-Containing Trisaccharide Related to Insulin Signal Transduction

Christine H. Jaworek, Pericles Calias,
Sarah Iacobucci, and Marc d'Alarcao*

Michael Chemistry Laboratory, Department of Chemistry
Tufts University, Medford, MA 02155

Received 7 October 1998; revised 23 November 1998; accepted 25 November 1998


ABSTRACT

A synthesis of D-mannosyl-(α 1-4)-D-glucosaminyl-(α 1-6)-D-*myo*-inositol-1:2-cyclic phosphate (**1**) from D-xylose, D-mannose, and D-glucal is reported. Compound **1** constitutes the terminal structure of the PI-PLC-released VSG membrane anchor of Trypanosomes and is structurally related to the inositol phosphate glycan implicated as an insulin second messenger in higher organisms.

© 1999 Elsevier Science Ltd. All rights reserved.

For the last dozen years, evidence has been accumulating that inositol-containing glycans play an important role in signal transduction in various cells. It has been demonstrated that insulin stimulates the release of inositol phosphate glycans (IPG's) from the cell membrane of insulin-sensitive cells and that these soluble molecules possess insulin-like properties.¹ Similar experiments have suggested that interleukin-1,² interleukin-2,³ and nerve growth factor⁴ also result in IPG release in their target cells. However, considerable structural diversity of these IPG molecules has been reported, including the presence of D-*chiro*-inositol and D-galactosamine⁵ vs. *myo*-inositol and D-glucosamine,⁶ and D-galactose⁷ vs. D-mannose.⁵ As yet, no definitive relationship between structure and signalling capability has been established for the IPG's.⁸ Accordingly, preparation of biologically active IPG's by unambiguous chemical synthesis is desirable to deconvolute the structural subtleties and elucidate the signalling pathway(s).

It has been established that the pentasaccharide VSG membrane anchor **2** derived from *Trypanosoma brucei* (v118) by Pronase treatment has significant insulin-like activities in several intact cell assays,⁹ probably due to its structural similarity with the IPG second messenger(s). It has also been found that synthetic *myo*-inositol-containing disaccharide cyclic phosphate **3**,¹⁰ and related *chiro*-inositol-containing disaccharide phosphates¹¹ exhibit limited insulin-like activity. Accordingly, information regarding the biological activity of molecules spanning this structural range (i.e. di- to penta-saccharides) is of greatest interest. In this communication, we describe the chemical synthesis of D-mannosyl-(α 1-4)-D-glucosaminyl-(α 1-6)-D-*myo*-inositol-1:2-cyclic phosphate (**1**). This compound constitutes the terminal trisaccharide portion of the insulin-mimetic VSG membrane anchor **2** from *T.brucei*.



A suitably protected *myo*-inositol derivative was prepared from optically pure inositol **4**, available from D-xylose via a known sequence.¹² Treatment of **4** with N, N'-carbonyldiimidazole followed by removal of the silyl protecting group afforded the differentially protected inositol **5**,¹³ suitable for glycosylation.


The glucosamine precursor, **6**, was prepared from tri-*O*-acetyl-D-glucal by the known procedure.¹⁴ Glycosylation of **6** with methylthio mannoside **7**,¹⁵ according to Ogawa's method,¹⁶ provided α -disaccharide **8**¹³ in 53% yield. Compound **8** was subjected to fluoride-promoted desilylation, converted to the glycosyl fluoride (DAST), and coupled with inositol **5** via Cp₂ZrCl₂-AgOTf activation¹⁷ to produce trisaccharide **9**.¹³

Selective removal of the cyclic carbonate moiety of **9** followed by phosphorylation of the resulting diol with the reagent generated from MeOPOCl₂ and pyridine¹⁸ produced the fully protected trisaccharide cyclic phosphate. Finally, removal of all benzyl protecting groups and reduction of the azido group was accomplished by treatment with sodium in liquid ammonia. The dissolving metal reduction reaction was quenched with NH₄Cl (s) at -78 °C followed by addition of MeOH at -78 °C and warmed to 20 °C. Desalting of **1** via Sephadex G-10 column chromatography (H₂O) yielded zwitterion **1**¹³ (56% over 2 steps).

The biological properties of trisaccharide **1** and related compounds are currently being evaluated.

Acknowledgment

We are grateful to the National Institutes of Health (grant # DK44589) for financial support of this work.



(a) *N,N'*-carbonyldiimidazole, benzene, 20 °C, 17h; (b) TBAF, THF, 0 °C, 3 min (84% over 2 steps); (c) PhSeCl, AgOTf, toluene, -42 °C, 1 h (53%); (d) *n*-BuN₄F, HOAc, THF, 20 °C (84%); (e) DAST, THF, -42 °C, 30 min (90%); (f) **5**, Cp₂ZrCl₂, AgOTf, toluene, -42 °C to 20 °C, 5h (28% α; 26% β); (g) 1 M LiOH, THF, 20 °C, 12h, (80%); (h) MeOPOCl₂, pyridine; (i) Na, NH₃ (l), -78 °C, 15 min (56% over 2 steps).

References

1. Saltiel, A. R.; Cuatrecasas, P. *Am. J. Physiol.* **1988**, *255*, C1.
2. Dobson, P. R. M. and Brown, B. L. In *The Biology and Medicine of Signal Transduction*; Nishizuka, Y. *et al.*, Eds.; Raven Press: New York; 1990; pp.170-175.
3. Merida, I.; Pratt, J. C.; Gaulton, G. N. *Proc. Natl. Acad. Sci. USA* **1990**, *87*, 9421-9425. Early, D. D.; Koshland, M. E. *Science* **1991**, *251*, 78-81.
4. Represa, J.; Avila, M. A.; Miner, C.; Giraldez, F.; Romero, G.; Clemente, R.; Mato, J. M.; Varela-Nieto, I. *Proc. Natl. Acad. Sci. USA* **1991**, *88*, 8016-8019.
5. Larner, J.; Huang, L. C.; Schwartz, C. F. W.; Oswald, A. S.; Shen, T. Y.; Kinter, M.; Tang, G.; Zeller, K. *Biochem. Biophys. Res. Commun.* **1988**, *151*, 1416.
6. Saltiel, A. R.; Cuatrecasas, P. *Proc. Natl. Acad. Sci. USA* **1986**, *83*, 5793.
7. Mato, J. M.; Kelly, K. L.; Abler, A.; Jarett, L.; Corkey, B.; Cashel, J.A.; Zopf, D. *Biochem. Biophys. Res. Commun.* **1987**, *146*, 764.

8. A recent paper reports considerable progress in this regard: Frick, W.; Bauer, A.; Bauer, J.; Wied, S.; Muller, G. *Biochemistry* **1998**, *38*, 13421-36.
9. Misek, D. E.; Saltiel, A. R. *J. Biol. Chem.* **1992**, *267*, 16266.
10. Plourde, R.; d'Alarcao, M.; Saltiel, A. R. *J. Org. Chem.* **1992**, *57*, 2606; Zapata, A.; León, Y.; Mato, J. M.; Varela-Nieto, I.; Penadés, S.; Martín-Lomas, M. *Carbohydr. Res.* **1994**, *264*, 21. Garegg, P.J. Konradsson, P.; Oscarson, S.; Ruda, K. *Tetrahedron* **1997**, *53*, 17727.
11. Reddy, K. K.; Falck, J. R.; Capdevila, J. *Tetrahedron Lett.* **1993**, *34*, 7869.
12. Kornienko, A.; Turner, D. I.; Jaworek, C. H.; d'Alarcao, M. *Tetrahedron: Asymmetry* **1998**, *9*, 2783-6.
13. Spectral data for **5**: ^1H NMR (CDCl_3) δ 3.88 (dd, $J=4.5, 10.7$ Hz, 1H), 3.80 (dd, $J=2.4, 4.5$ Hz, 1H), 3.90 (wt, $J=2.8$ Hz, 1H), 4.25-4.75 (m, 8H), 4.85 (dd, $J=3.3, 8.7$ Hz, 1H), 7.15-7.5 (m, 15H, phenyl). **8**: ^1H NMR (CDCl_3) δ 0.19 (s, 3H, CH_3), 0.20 (s, 3H, CH_3), 0.94 (s, 9H, $t\text{-Bu}$), 3.25 (dd, $J=8.6, 9.9$ Hz, 1H, H3 gln), 3.36-3.42 (m, 2H, H5 & H2 gln), 3.58 (dd, $J=1.5, 9.1$ Hz, 1H, H3 man), 3.64-3.83 (m, 7H), 3.98 (wt, $J=9.3$ Hz, 1H, H4 man), 4.25 (d, $J=12.2$ Hz, 1H, 1/2 CH_2Ph), 4.35 (d, $J=12.2$ Hz, 1H, 1/2 CH_2Ph), 4.43-4.63 (m, 9H), 4.82 (d, $J=10.8$ Hz, 1H, 1/2 CH_2Ph), 4.94 (d, $J=11.4$ Hz, 1H, 1/2 CH_2Ph), 5.30 (d, $J=2.0$ Hz, 1H, H1 man), 7.14-7.40 (m, 30H, Ph). ^{13}C NMR (CDCl_3) δ 100.15 (d, $J=169$ Hz, C1 man), 97.15 (d, $J=160$ Hz, C1 gln). **9**: ^1H NMR (CDCl_3) δ 3.39 (dd, $J=3.6, 9.7$ Hz, 1H, H2 gln), 3.51-3.90 (m, 12H), 4.02 (wt, 2H), 4.20-4.66 (m, 17H), 4.79-5.02 (m, 4H), 5.27 (d, $J=1.9$ Hz, 1H, H1 man), 5.34 (d, $J=3.6$ Hz, 1H, H1 gln), 7-7.5 (45H, m, Ph). LR FAB (pos. ion mode): Found: 1388; Calculated for $\text{C}_{82}\text{H}_{83}\text{N}_3\text{O}_{16} + \text{Na}$: 1388.57. **1**: ^1H NMR (D_2O) δ 3-4 (m, 16H), 4.35 (ddd, 1H, H1 inos), 4.5 (wt, 1H, H2 inos), 5.1 (vs, 1H), 5.3 (vs, 1H). ^{31}P NMR (D_2O): 17.135 ppm (relative to 85% H_3PO_4). HR FAB MS (negative ion mode): Found: 564.1354; Calculated for $\text{C}_{18}\text{H}_{32}\text{NO}_{17}\text{P} - \text{H}$: 564.1329.
14. Murakata, C.; Ogawa, T. *Carbohydr. Res.* **1992**, *234*, 75.
15. Compound **7** was prepared from 2,3,4,6-tetra-*O*-benzyl-D-mannopyranose by fluorination with DAST followed by treatment with Bu_4SnSMe and SnCl_4 in overall 89% yield (α : β , 14:1). See: Yamazaki, F.; Sato, S.; Nukada, T.; Ito, Y.; Ogawa, T. *Carbohydr. Res.* **1990**, *201*, 31.
16. Ito, Y.; Ogawa, T. *Tetrahedron Lett.* **1988**, *29*, 1061.
17. Matsumoto, T.; Maeta, H.; Suzuki, K.; Tsuchihashi, G. *Tetrahedron Lett.* **1988**, *29*, 3567.
18. Smrt, J.; Catlin, J. *Tetrahedron Lett.* **1970**, 5081; Wantanabe, Y.; Ogasawara, T.; Nakahira, H.; Matsuki, T.; Ozaki, S. *Tetrahedron Lett.* **1988**, *29*, 5259.